

Counting Marbles With
'Accessible' Mass Density:
A Reply to Bassi and Ghirardi

Rob Clifton

*Departments of Philosophy and History and Philosophy of Science,
University of Pittsburgh, Pittsburgh, PA 15260
(e-mail: rclifton+@pitt.edu)*

and

Bradley Monton

*Department of Philosophy, 1879 Hall, Princeton University, Princeton, NJ
08544-1006
(e-mail: bjmonton@princeton.edu)*

February 1, 2008

1 Review of the Counting Anomaly Debate

In the previous issue of this journal ([1999]) we argued that, while Lewis ([1997]) is correct that the enumeration principle fails in dynamical reduction theories, one need not follow Lewis in rejecting these theories. Because the dynamical reduction process itself prevents the failure of enumeration from ever becoming manifest, and because one can treat the semantics for dynamical reduction theories as not adding anything of ontological import to them, it is reasonable to accept these theories notwithstanding Lewis's counting anomaly.

Lewis had considered a collection of n non-interacting marbles in the product state

$$|\psi\rangle_{\text{all}} = (a|\text{in}\rangle_1 + b|\text{out}\rangle_1) \otimes (a|\text{in}\rangle_2 + b|\text{out}\rangle_2) \otimes \cdots \otimes (a|\text{in}\rangle_n + b|\text{out}\rangle_n), \quad (1)$$

where the state $|\text{in}\rangle_i$ refers to localization of the i th marble to inside a box (assumed, for simplicity, to be the same box for all marbles), and $1 > |a|^2 \gg |b|^2 > 0$. These inequalities are strict because dynamical reduction theories generically do not eliminate the tails of an object's wavefunction when it collapses. Nevertheless, since $|a|^2 \gg |b|^2$, the probability of finding any particular marble in the box is very high, and it is natural to take each individual marble to be in the box. Indeed, not doing so would leave dynamical reduction theories without a complete solution to the measurement problem (see also Albert and Loewer [1996]). On the other hand, for sufficiently large n , the probability that *all* n marbles will be found in the box is $|a|^{2n} \ll 1$, making it highly unlikely that they will all be found in the box. Lewis inferred from this that the state $|\psi\rangle_{\text{all}}$ 'cannot be one in which all n marbles are in the box, since there is almost no chance that if one looks one will find them there' ([1997], p. 318). We cashed out Lewis's inference by viewing it as a consequence of what we called the *fuzzy* eigenstate-eigenvalue link:

'Particle x lies in region R_x and y lies in R_y and z lies in R_z and

...’ if and only if the proportion of the total squared amplitude of $\psi(t, \mathbf{r}_1, \dots, \mathbf{r}_N)$ that is associated with points in $R_x \times R_y \times R_z \times \dots$ is greater than or equal to $1 - p$.

The essence of Lewis’s counting anomaly, then, is that the application of the fuzzy link in state $|\psi\rangle_{\text{all}}$ to the collection of marbles as a whole conflicts with its application on a marble-by-marble basis. And the only way out of this difficulty is to abandon an instance of the logical rule of conjunction introduction that Lewis called the *enumeration principle*: ‘if marble 1 is in the box and marble 2 is in the box and so on through marble n , then all n marbles are in the box’ ([1997], p. 321).

Since we were defending this part of Lewis’s argument, we were obliged to evaluate counter-arguments to it. Thus, in section 3 of our paper, we criticized three arguments against Lewis’s claim, the main one being due to Ghirardi and Bassi ([1999]). In their response to our paper, Bassi and Ghirardi ([1999]) (whom we shall henceforth refer to simply as ‘BG’) reject our criticism, as well as our argument against Lewis that dynamical reduction precludes failures of conjunction introduction from ever becoming manifest. Our intention here is to demonstrate that BG’s responses to us do not succeed.

2 The Counting Anomaly Dies Hard

Ghirardi and Bassi’s ([1999]) rejection of Lewis’s counting anomaly was based, in part, upon calculating that, for an object the size of a marble, the coefficient a occurring in $|\psi\rangle_{\text{all}}$ will turn out to be so close in absolute square to 1 that, even if all the mass in the actual universe were used to form the marbles, n would not be large enough for $|a|^{2n}$ to be appreciably smaller than 1. In our reply ([1999], pp. 17-8), we pointed out that one could equally well pose the anomaly in terms of counting sufficiently many *microscopic* objects, such as elementary particles, whose a coefficients would

certainly *not* be forced by dynamical reduction to be exceedingly close to 1. Clearly it would be no less strange if arithmetic failed for the particles out of which things like marbles are made up! Moreover, the virtue of our own response to the anomaly (i.e., that it is never manifest) is that it applies in any world governed by dynamical reduction, independent of contingent limits on marble or particle aggregation in our world. This leaves us puzzled about why BG continue to emphasize that the magnitude of the number n required to get the counting anomaly off the ground is ‘unphysically large’ ([1999], p. 15), and that it is a ‘peculiar game of putting no limit on the number of marbles’ ([1999], p. 16). We echo their sentiment that ‘we do not like to play with science fiction arguments’ ([1999], p. 16), but it is not as if one is being asked to consider an *actual infinity* of objects.

In any case, Ghirardi and Bassi’s ([1999]) main argument against the counting anomaly was based upon rewriting the state $|\psi\rangle_{\text{all}}$ as

$$\begin{aligned} |\psi\rangle_{\text{all}} = & a^n |\text{in}\rangle_1 \otimes |\text{in}\rangle_2 \cdots |\text{in}\rangle_n + a^{n-1} b |\text{out}\rangle_1 \otimes |\text{in}\rangle_2 \cdots |\text{in}\rangle_n \\ & + a^{n-1} b |\text{in}\rangle_1 \otimes |\text{out}\rangle_2 \cdots |\text{in}\rangle_n + \cdots + a^{n-1} b |\text{in}\rangle_1 \otimes |\text{in}\rangle_2 \cdots |\text{out}\rangle_n \\ & + a^{n-2} b^2 |\text{out}\rangle_1 \otimes |\text{out}\rangle_2 \cdots |\text{in}\rangle_n + \cdots + b^n |\text{out}\rangle_1 \otimes |\text{out}\rangle_2 \cdots |\text{out}\rangle_n, \end{aligned} \quad (2)$$

and then asserting: ‘the precise GRW dynamics will lead in about one millionth of a second to the suppression of the superposition and to the “spontaneous reduction” of the state [(2)] to one of its terms’ ([1999], p. 52). Were such a reduction to occur, conjunction introduction would certainly hold for the resultant state. Therefore, Ghirardi and Bassi concluded that conjunction introduction holds, since a state like (2) ‘never occurs, its existence being forbidden by the GRW theory itself’ ([1999], p. 52). In our paper, however, we argued that Ghirardi and Bassi’s characterization of the reduction of the state (2) was incorrect and did nothing to remove the anomaly. While BG implicitly concede the error ([1999], pp. 10-11), they now claim:

... the point we wanted to make is that a rigorous interpretation of states of this type requires a consideration of the mass density

functional ... and cannot be based on the *fuzzy link* criterion or other inappropriate criteria. Therefore, the criticism of Clifton and Monton ([1999]) to our previous paper misses the crucial point of our argument just because it does not make reference to valid criteria for interpreting the wave function within dynamical reduction models ([1999], p. 11).

We definitely did not see that the crucial point of Ghirardi and Bassi's ([1999]) argument against Lewis was a rejection of fuzzy link-style semantics in favor of the mass density interpretation. We think we can be forgiven for the oversight given that reference to the mass density interpretation in their ([1999]) only occurs in three places — two footnotes and a parenthesis — and, at the second occurrence (footnote 3), they state that their arguments against Lewis could equally well be made under the mass density interpretation (which has the effect of suggesting that their arguments did not critically depend upon that interpretation). Moreover, while it may now be the case that the mass density approach is 'universally accepted' by dynamical reduction theorists (BG [1999], p. 7), one can find explicit endorsement in Pearle ([1997], pp. 150-1) of fuzzy link-type semantics applied to quantities such as local charge, spin, energy, and momentum (in particular, see equation (11) of Pearle (1997)). The important issue, of course, is whether the counting anomaly arises under the mass density interpretation (setting aside, for the next section, the issue of whether the anomaly can be made manifest). We shall argue that either the counting anomaly arises, or some other (equally surprising) anomaly will arise.

We start by reviewing a nice example that Ghirardi, Grassi, and Benatti ([1995]) use to first introduce the mass density interpretation. Consider two disjoint spherical regions of space A and B , each large enough to contain a macroscopic number N of particles, each of mass m . Let $|\phi_N^A\rangle$ denote a state where the N particles are each 'well-localized with respect to the characteristic length (10^{-5} cm) of the model and uniformly distributed' in

region A ([1995], p. 16), and consider two possible states of the N -particle system:

$$|\psi^+\rangle = 1/\sqrt{2}(|\phi_N^A\rangle + |\phi_N^B\rangle) \quad , \quad |\psi^\otimes\rangle = |\phi_{N/2}^A\rangle \otimes |\phi_{N/2}^B\rangle. \quad (3)$$

For simplicity, Ghirardi *et al* divide space up into cells with dimensions given by the characteristic length. They then consider the expectation of the mass operator M_i for a cell i ,

$$\mathcal{M}_i = \langle \psi | M_i | \psi \rangle. \quad (4)$$

It is easy to see that both $|\psi^+\rangle$ and $|\psi^\otimes\rangle$ give rise to the same mass density function for a cell i in region A :

$$\mathcal{M}_i^+ = \langle \psi^+ | M_i | \psi^+ \rangle \approx 1/2 \langle \phi_N^A | M_i | \phi_N^A \rangle \approx nm/2, \quad (5)$$

$$\mathcal{M}_i^\otimes = \langle \psi^\otimes | M_i | \psi^\otimes \rangle \approx \langle \phi_{N/2}^A | M_i | \phi_{N/2}^A \rangle \approx nm/2, \quad (6)$$

where n is the number of particles per cell. However, Ghirardi *et al* argue that only one of these masses is objective. They consider what happens if one sends a test mass between regions A and B . If the system were in state $|\psi^\otimes\rangle$, gravitational forces would balance and the particle would be undeflected. On the other hand, if the system were in state $|\psi^+\rangle$, the test particle would become involved in the superposition, where in one branch it is deflected toward A and in the other toward B . Ghirardi *et al* write:

Nowhere in the universe is there a density corresponding to the density of the test particle. In a sense, if one would insist in giving a meaning to the density function, he would be led to conclude that the particle has been split by the interaction into two pieces of half its density. This analysis shows that great attention should be paid in attributing an ‘objective’ status to the function $\mathcal{M}(\mathbf{r})$ ([1995], p. 17; see also Ghirardi and Grassi [1996], p. 365).

Ghirardi *et al* are then led to propose that the mass in a cell i is ‘objective’ just in case $\mathcal{R}_i \equiv \mathcal{V}_i / \mathcal{M}_i^2 \ll 1$, where the variance of M_i is defined in the

usual way by

$$\mathcal{V}_i = \langle \psi | [M_i - \langle \psi | M_i | \psi \rangle]^2 | \psi \rangle. \quad (7)$$

Only in state $|\psi^\otimes\rangle$ is $\mathcal{R}_i \ll 1$ for a cell in A ; for the state $|\psi^+\rangle$, one has $\mathcal{R}_i \approx 1$.

Note that $\mathcal{R}_i \ll 1$ implies that any mass measurement in cell i is almost certain to yield $mn/2$ as its answer. In fact, this is what BG ([1999]) take to be the main motivation for the proposed objective mass density criterion; for, in their own introduction to the criterion, they reproduce the following passage from Ghirardi ([1997]):

A property corresponding to a value (or range of values) of a certain variable in a given theory is objectively possessed or accessible when, according to the predictions of that theory, experiments (or physical processes) yielding reliable information about the variable would, if performed (or taking place), give an outcome corresponding to the claimed value. Thus the crucial feature characterizing accessibility (as far as statements of individual systems is concerned) is the matching of the claims and the outcomes of physical processes testing the claims (Ghirardi [1997], p. 227).

Surely if these remarks serve to motivate the mass density criterion, they also serve to motivate speaking of a particle (or particles) as being located in a region whenever its wavefunction assigns high probability to its being detected in that region; that is, they also serve to motivate the fuzzy link. Unfortunately, BG never say why the fuzzy link is ‘inappropriate’ and not a ‘valid’ way to understand reduction theories ([1999], p. 11)¹.

¹One might surmise that the fuzzy link, which is formulated in terms of configuration, is unacceptable to BG because there is no well-behaved relativistic configuration operator (e.g., see Malament [1996]). However, it does not matter whether one conducts discussion of the counting anomaly within a nonrelativistic or relativistic setting. If one wants to insist on the latter, then it is trivial to reformulate the fuzzy-link in terms of observables that *are* relativistically acceptable (local charge, energy, etc.).

It is also unclear, when BG invoke the ‘physically more expressive term’ *accessible* ([1999], p. 7), just how seriously we are to take talk of mass density as ‘objective’ when the condition $\mathcal{R}_i \ll 1$ obtains. Elsewhere, Ghirardi and Grassi ([1996], p. 376) have written, with regard to the term ‘objective’, that ‘both usual meanings of that term (i.e. “real” or “opposite to subjective”) do not fit with the sense which emerges for it from our work’. Our response to BG’s criticism of the counting anomaly based on mass density will not in any way depend on how these terms are to be understood. However, it is worth reminding the reader of the tentative position that we arrived at in our ([1999]): that the fundamental ontology of dynamical reduction theories consists only of wavefunctions and their evolution, and that a fuzzy link or mass density semantics, though essential for connecting the theory to our everyday experience and conception of the world, need not be seen as essential for making ontological sense of such theories.

We return, now, to the counting anomaly. In order for mass density to be relevant to the issue of how many particles are in the box, we need a way to connect mass density talk to position talk. Ghirardi *et al* already mention the obvious way of doing this: ‘The reality of a massive macro-object in front of us corresponds to the fact that in the region it occupies there is the objective mass density which characterizes it’ ([1995], p. 33). And this is BG’s understanding of things, when they assert:

... the mass is objective precisely in the regions where the various marbles are... The marbles can therefore be claimed to be all within the box, and the total mass within the box is actually the one corresponding to all of them being in the box ([1999], p. 9).

It is the second sentence with which we disagree. Consider, first, a single marble in state $a|\text{in}\rangle + b|\text{out}\rangle$, where $|a| \gg |b|$. If the marble’s total mass, i.e., mass density function integrated over all space, is m , then the total accessible mass in the box will only be $|a|^2 m$, and this can of course change with time. If one thought that an essential property of a marble is that

its objective (rest) mass is constant, one might resist locating the marble entirely in the box. We are not inclined to resist that ourselves, since the task, as we see it, is to supply a servicable mapping between the theory and our language about marbles. However, the plot thickens when one considers an n -marble system in state $|\psi\rangle_{\text{all}}$. In that case, Lewis ([1997], appendix A) provided a detailed analysis, after which he arrived at the same conclusion as BG (!):

There are n regions of objective mass $|a|^2 m$ in the box, one corresponding to each marble, and no regions outside the box; consequently the mass distribution is such that all n marbles are in the box (1997, p. 327).

Unfortunately, we fail to see how Lewis's conclusion here follows. Since the total mass in the box is $(|a|^2 m) \times n$, n can be chosen sufficiently large so that total does not exceed, for example, $m(n - 1000)$. Since one could well believe that it is inappropriate to assert that all n marbles are in the box unless roughly mn mass is objective therein, conjunction introduction need not be satisfied. And it is little help to respond that, for n sufficiently large, the number $m(n - 1000)$ is roughly the same as mn . For one could still be puzzled as to why it would not then be equally legitimate to assert that $n - 1000$ marbles are in the box as well.

It should be emphasized that even though Lewis claimed not to find any failure of conjunction introduction under the mass density interpretation, he maintained that it was unacceptable to assert that all marbles are in the box, on the grounds that there is still a vanishing probability in state $|\psi\rangle_{\text{all}}$ of finding them all there (1997, p. 320). Though BG are dismissive of Lewis on this point ([1999], note 2), we entirely agree with him (and gave supporting arguments for his position in our [1999], pp. 21-2). Note also that if one wants to maintain that all the mass of the marbles is objectively in the box, together with the fact that the probability of finding them all there is vanishingly small, then one is committed to a radical breach between

mass and location talk. Thus we can pose BG a trilemma for the mass density interpretation: either conjunction introduction fails, mass talk must be divorced from position talk, or the intuitive connection between either of these kinds of talk and a system's dispositions (cf. 'accessibility') must be severed.

3 Suppressing the Counting Anomaly

In our ([1999]), we defended the fuzzy link in the face of Lewis's counting anomaly by showing that a violation of conjunction introduction could never be made manifest. Notwithstanding BG's critique, we stand by our arguments. Indeed, we believe that they apply equal well in defence of the mass density criterion!

Our escape from the counting anomaly was based on Lewis's failure to explicitly model, or 'operationalize', the counting process itself within dynamical reduction theories. BG begin their critique by rejecting this line of thinking out of hand:

The unappropriateness of such a point of view has already been stressed in section 3. Within GRW's theory no *measurement* process ever occurs, the only physical processes being interactions among physical systems governed by universal laws ([1999], p. 17).

Thus, it is worth being clear that nothing in our analysis turned on privileging (so-called) *measurement* interactions.

In that analysis, we considered what would happen when devices M_i , each designed to register whether or not marble i is in the box, are coupled to the marbles in state $|\psi\rangle_{\text{all}}$; and when a system M , designed to register the *total* number of particles in the box, is coupled to the total n -marble system. Just after interaction with all the registers, the state would be (using an obvious

notation):

$$\begin{aligned} |\psi\rangle_{\text{count}} &= a^n |\phi\rangle_{\text{out}_0} |O = n\rangle_M + a^{n-1} b |\phi\rangle_{\text{out}_1} |O = n-1\rangle_M \\ &\quad + \cdots + b^n |\phi\rangle_{\text{out}_n} |O = 0\rangle_M, \end{aligned} \quad (8)$$

where

$$\begin{aligned} |\phi\rangle_{\text{out}_0} &= |\text{in}\rangle_1 |\text{'in'}\rangle_{M_1} |\text{in}\rangle_2 |\text{'in'}\rangle_{M_2} \cdots |\text{in}\rangle_n |\text{'in'}\rangle_{M_n} \\ |\phi\rangle_{\text{out}_1} &= |\text{out}\rangle_1 |\text{'out'}\rangle_{M_1} |\text{in}\rangle_1 |\text{'in'}\rangle_{M_2} \cdots |\text{in}\rangle_n |\text{'in'}\rangle_{M_n} \\ &\quad + |\text{in}\rangle_1 |\text{'in'}\rangle_{M_1} |\text{out}\rangle_1 |\text{'out'}\rangle_{M_2} \cdots |\text{in}\rangle_n |\text{'in'}\rangle_{M_n} \\ &\quad \vdots \\ &\quad + |\text{in}\rangle_1 |\text{'in'}\rangle_{M_1} |\text{in}\rangle_1 |\text{'in'}\rangle_{M_2} \cdots |\text{out}\rangle_n |\text{'out'}\rangle_{M_n} \\ &\quad \vdots \\ |\phi\rangle_{\text{out}_n} &= |\text{out}\rangle_1 |\text{'out'}\rangle_{M_1} |\text{out}\rangle_2 |\text{'out'}\rangle_{M_2} \cdots |\text{out}\rangle_n |\text{'out'}\rangle_{M_n}, \end{aligned}$$

We then argued that dynamical reduction will rapidly reduce the entangled state $|\psi\rangle_{\text{count}}$ to one in which the individual recordings of the n registers always agree with whatever total M records. The argument depends only the fact that, *unlike* $|\psi\rangle_{\text{all}}$, $|\psi\rangle_{\text{count}}$ is an *entangled* state involving a macroscopic number of particles. Moreover, one need not assume that M , or any of the M_i , are themselves macroscopic (cf. our [1999], notes 6,7). Of course, in order for the marble count(s) to be *known* by the ‘gross creatures that we are’ (to echo a phrase of John Bell’s), these devices would have to be coupled to macroscopic devices. But the suppression of the counting anomaly will have occurred long before information about the marbles ever enters anyone’s consciousness — it is the purely physical outcome of purely physical interactions and correlations.

Criticizing our discussion of the various possible collapse scenarios that could follow interaction with the registers, BG ask rhetorically:

The marbles are macroscopic systems and their masses may very well be comparable or even larger than those of the pointers both

of the apparatuses M_i and of M . Why the GRW dynamics has to be suspended up to the time in which the pointer of M is localized, and subsequently (in the case that $O \neq n$) up to the moment in which all pointers of the M_i 's are localized, is really a big mystery ([1999], pp. 20-1).

In our exposition, we were simply dividing the collapse process into different stages for ease of exposition, not trying to give a real-time description of the dynamics. Certainly we were aware that the marbles themselves will be almost continually subject to GRW collapses before, during, and after the interaction with the registers, causing marbles to jump in and out of the box almost all the time (provided enough marbles are considered). As BG say, ‘The situation...changes every millionth of a second’ ([1999], p. 16). However, we fail to see why this means we:

...must first of all assume that it is possible to perform the incredible task of testing the positions of all particles and the total mass within the box in much less than a millionth of a second ([1999], p. 16).

For, as we argued ([1999], pp. 18-9), at *every* moment prior to the entangling interaction with M , there will always be some conjunctive assertion about the marbles’ locations relative to the box with respect to which conjunction introduction fails. The point of introducing M was not so it can provide a faithful recording of the total marble count *prior* to its interaction with the marbles, but to demonstrate that the *most current* marble count will always agree with the *most current* information contained in the M_i registers.

BG summarize our response to Lewis with the phrase: ‘the violation of the counting rule is still there, but it cannot be revealed’ ([1999], p. 20). This is not correct: the marbles’ violation of the enumeration principle will itself disappear, along with any chance of *manifesting* such a violation, a split second after their interaction with M (i.e., as soon as $|\psi\rangle_{\text{count}}$ collapses).

Maintaining their resistance to this conclusion, BG finish their critique by apparently arguing that a failure of conjunction introduction in the marbles system could, after all, be made manifest:

Since M 's reading is given by its pointer location . . . a consistent use of the GRW dynamics implies that *it cannot* be perfectly localized: it will be affected by the 'tails problem' just as all macroscopic systems. Thus, if one follows the authors in their strange argument, one should conclude that (with a small probability) the state can be the one in which n marbles are in the state $|\text{in}\rangle$, n apparatuses M_i register $|\text{'in'}\rangle$, but the pointer of the apparatus M can be reduced on a state corresponding to its pointing at $k \neq n$! ([1999], p. 21)

In our analysis, we ignored the subsequent development of tails in the wavefunction of M 's pointer after its interaction with the marbles, and treated that interaction as 'perfect' in the sense that, if the marbles had been in an eigenstate of a definite number of particles, j , in the box, M would register $O = j$. Dropping these idealizations would simply mean that M has some small probability of registering an incorrect result for the total marble count. But that kind of measurement error is strictly irrelevant to manifesting a failure of enumeration. Indeed, our point was that even if one assumes that the registers operate as perfectly as could be, they will not display any counting anomaly. As we put it in the paper (in relation to an observer playing the role of M): 'the only way to arrange things so that our observer *could* falsify the enumeration principle would be to suppose that she was never a competent enumerator to begin with!' ([1999], p. 31).

References

Albert, D. and Loewer, B. [1996]: 'Tails of Schrödinger's Cat', in R. Clifton (ed), *Perspectives on Quantum Reality*, Dordrecht: Kluwer, pp. 81-91.

- Bassi, A., and G. C. Ghirardi [1999]: ‘More About Dynamical Reduction and the Enumeration Principle’, *The British Journal for Philosophy of Science*, December issue.
- Clifton, R. and B. Monton [1999]: ‘Losing Your Marbles in Wavefunction Collapse Theories’, *The British Journal for Philosophy of Science*, December issue.
- Ghirardi, G. C. [1997]: ‘Macroscopic Reality and the Dynamical Reduction Program’, in M. L. Dalla Chiara *et al* (eds), *Structures and Norms in Science*, Dordrecht: Kluwer, pp. 221-240.
- Ghirardi, G. C. and A. Bassi [1999]: ‘Do Dynamical Reduction Models Imply That Arithmetic Does Not Apply to Ordinary Macroscopic Objects?’, *The British Journal for Philosophy of Science*, **50**, pp. 705-720.
- Ghirardi, G. C. and R. Grassi [1996]: ‘Bohm’s Theory versus Dynamical Reduction’, in J. T. Cushing *et al* (eds), *Bohmian Mechanics and Quantum Theory: An Appraisal*, Dordrecht: Kluwer, pp. 353-377.
- Ghirardi, G. C., R. Grassi, and F. Benatti [1995]: ‘Describing the Macroscopic World: Closing the Circle within the Dynamical Reduction Program’, *Foundations of Physics*, **25**, pp. 5-38.
- Lewis, P. [1997]: ‘Quantum Mechanics, Orthogonality, and Counting’, *The British Journal for Philosophy of Science*, **48**, pp. 313-328.
- Malament, D. [1996]: ‘In Defense of Dogma: Why There Cannot be a Relativistic Quantum Mechanics of (Localizable) Particles’, in R. Clifton (*ed*), *Perspectives on Quantum Reality*, Boston: Kluwer, pp. 1-10.
- Pearle, P. [1997]: ‘Tales and Tails and Stuff and Nonsense’, in R. S. Cohen, M. Horne, and J. Stachel (*eds*), *Experimental Metaphysics*, Dordrecht: Kluwer, pp. 143-56.